



# Gentaur

## E-cadherin Stable MIA PaCa-2 Cell Line

<b>Cat.No.</b>	<b>Unit</b>
T3167	1x10 <sup>6</sup> cells / 1.0 ml

---

<b>Cat. No.</b>	T3167
<b>Name</b>	E-cadherin Stable MIA PaCa-2 Cell Line
<b>Description</b>	<p>E-cadherin (Ecad) is prominently studied for its role in tumorigenesis due the loss of E-cadherin is a critical initiation step in the epithelial-to-mesenchymal transition associated with the invasiveness of tumors causing poor prognosis. The E-cadherin Stable MIA PaCa-2 Cell Line is a tool for tumor-related studies on the effects Ecad has in cancer prognosis and antitumor therapies. Ecad is a relevant biomarker for therapies, such as for TK/GCV (thymidine kinase/ganciclovir) therapy.</p>
<b>Organism</b>	Human (H. sapiens)
<b>Tissue</b>	Pancreas
<b>Donor History</b>	Male, 65, Caucasian, Carcinoma
<b>Growth Properties</b>	Adherent, epithelial-like
<b>Cell Type</b>	Stable Cell Lines
<b>Unit</b>	1x10 <sup>6</sup> cells / 1.0 ml
<b>Storage Condition</b>	Vapor phase of liquid nitrogen, or below -130°C.
<b>Shipping Conditions</b>	Ship with dry ice.
<b>Product Format</b>	Frozen
<b>Intended Use</b>	This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.
<b>BioSafety</b>	II

<b>Certificate of Analysis</b>	For batch-specific test results, refer to the applicable certificate of analysis that can be found at <a href="http://www.abmgood.com">www.abmgood.com</a> .
<b>Growth Conditions</b>	Use of PriCoat™ T25 Flasks (G299) or Applied Cell Extracellular Matrix (G422) is required for cell adhesion to the culture vessels. PriGrow III (TM003) + 10% FBS + 2 mM L-glutamine + 1% Penicillin/Streptomycin Solution (G255), 37.0°C, 5% CO <sub>2</sub> . Selection with 0.5 - 2 µg/ml Puromycin (G264)
<b>Unpacking and Storage Instructions</b>	<ol style="list-style-type: none"> <li>1. Visually examine the packaging containers for signs of leakage or breakage.</li> <li>2. Immediately transfer frozen cells from dry ice packaging to a temperature below -130°C, preferably in liquid nitrogen vapor phase storage, until ready for use.</li> </ol> <p>To ensure the highest level of viability, thaw the vial and initiate culture as soon as possible upon receipt. If continued storage is desired, the vial should only be stored below -130°C or in liquid nitrogen vapor phase. Do not store at -70°C, as it will result in loss of viability.</p>
<b>Thawing Protocol</b>	<ol style="list-style-type: none"> <li>1. Thaw cells quickly in a 37°C water bath while agitating gently (maximum 2 minutes). The vial cap should be kept above the water level to minimize the risk of contamination.</li> <li>2. Decontaminate the vial by spraying and wiping the exterior of the vial with 70% ethanol. From this point onwards, all operations should be strictly carried out inside a biological safety cabinet using aseptic conditions.</li> <li>3. Transfer the cell suspension into a 15ml sterile conical tube containing 5ml of pre-warmed, complete growth media. Centrifuge cells at 125xg for 5-7 minutes.</li> <li>4. Aspirate the supernatant without disturbing the cell pellet. Re-suspend the cell pellet in the recommended pre-warmed, complete growth media and dispense into a T25 culture flask.</li> <li>5. Incubate the cells at the recommended conditions.</li> </ol>
<b>Subculture Protocol</b>	<p>Volumes given below are for a T75 flask; proportionally increase or decrease the volume as required per culture vessel size. Subculture cells once the culture vessel is 80% confluent.</p> <ol style="list-style-type: none"> <li>1. Aspirate the culture media, and add 2-3ml of pre-warmed 0.25% Trypsin-EDTA to the culture vessel.</li> <li>2. Observe the cells under a microscope to confirm detachment (typically within 2-10 minutes). Cells that are difficult to detach can be put in 37°C, for several minutes to facilitate detachment.</li> <li>3. Neutralize Trypsin-EDTA by adding an equal volume of the complete growth media into the culture vessel.</li> </ol>

4. Transfer the culture suspension into a sterile centrifuge tube, and centrifuge at 125xg for 5 minutes. The actual centrifuge duration and speed may vary depending on the cell type.
5. Aspirate the supernatant, and re-suspend the pellet with pre-warmed fresh complete growth media. Add appropriate aliquots of the cell suspension to new culture vessels, as desired.
6. Incubate the cells at the recommended conditions.

<b>Cryopreservation</b>	Cryopreservation Medium (TM024), or complete growth media with 10% DMSO.
<b>Expression</b>	E-cadherin
<b>Material Citation</b>	If use of this material results in a scientific publication, please cite the material in the following manner: Applied Biological Materials Inc, Cat. No. T3167.
<b>Warranty</b>	<b>abm</b> warrants that cell lines shall be viable upon initiation of culture for a period of thirty (30) days after shipment and that they shall meet the specifications on the applicable <b>abm</b> Material Product Information sheet, certificate of analysis, and/or catalog description. Such thirty (30) day period is referred to herein as the "Warranty Period".
<b>Disclaimer</b>	<ol style="list-style-type: none"><li>1. Sale of this item is subjected to the completion of a Material Transfer Agreement (MTA) by the purchasing individual/institution for each cell line. If you have any questions regarding this, please contact us at <a href="mailto:licensing@abmgood.com">licensing@abmgood.com</a>.</li><li>2. All test parameters provided in the CoA are conducted using <b>abm's</b> standardized culture system and The stated values may vary under the end-user's culture conditions. Please verify that the product is suitable for your studies by referencing published papers or ordering RNA (0.5 µg, Cat.# C207, \$450.00) or cell lysate (100 µg, Cat.# C206, \$600.00) to perform preliminary experiments, or alternatively use our Gene Expression Assay Service (Cat# C138). All sales are final.</li><li>3. We recommend live cell shipments for ease of cell transfer and this option can be requested at the time of order placement. Please note that the end-user will need to evaluate the feasibility of live cell shipment by taking into account the final destination's temperature variation and its geographical location.</li><li>4. All of <b>abm's</b> cell biology products are for research use ONLY and NOT for therapeutic/diagnostic applications. <b>abm</b> is not liable for any repercussions arising from the use of its cell biology product(s) in therapeutic/diagnostic or any other non-RUO application(s).</li><li>5. <b>abm</b> makes no warranties or representations as to the accuracy of the information on this site. Citations from literature are provided for informational</li></ol>

purposes only. **abm** does not warrant that such information has been shown to be accurate.

6. **abm** warrants that cell lines shall be viable upon initiation of culture for a period of thirty (30) days after shipment and that they shall meet the specifications on the applicable **abm** Material Product Information sheet, certificate of analysis, and/or catalog description. Such thirty (30) day period is referred to herein as the "Warranty Period".

**Depositor** Institut D'Investigacions Biomèdiques August Pi I Sunyer (IDIBAPS)  
**Application** Research Use Only.

---

**Gentaur Ltd.**

Unicorn House, 2nd floor, Station Close,  
Potters Bar, Hertfordshire, EN6 1TL, United Kingdom  
E: [uk@gentaur.com](mailto:uk@gentaur.com) | T: 020 33 93 85 31 | W: [www.gentaur.com](http://www.gentaur.com)